

OHSTPY-HEP-T-97-011
hep-ph/9707267

ON PERTURBATIVE QCD AT FINITE TEMPERATURE

A. NIETO

The Ohio State University, Department of Physics, Columbus, OH 43210^a

Effective field theory methods provide a convenient approach to study static observables in field theory at finite temperature. In this talk, I will outline the construction of the effective field theory that describes effective observables in QCD at high temperature. An analysis of the convergence of the perturbative series for the free energy of QCD will also be presented.

1 Introduction

There is a number of interesting scenarios which are described by quantum field theories at finite temperature (electroweak phase transition, heavy ion collisions, etc.) In this talk I will focus on the deconfinement phase transition of QCD: lattice simulations suggest that hadron matter undergoes a transition into a phase in which quarks are deconfined at a temperature of order 150 MeV. This new state of matter called *quark-gluon plasma (QGP)* is expected to be studied experimentally at CERN and Brookhaven in the next decade. Here, we will consider the system at high temperature so that it is well inside the QGP phase. The temperature is the typical energy of the system and, when it is high enough, the strong coupling constant is small so that one would expect perturbation theory to describe the QGP accurately.

It is specially interesting to consider the limit in which all of the masses scales of the system are smaller than the temperature. However, in this case, particles are massless and one has to deal with infrared divergences. In perturbation theory the infrared problems are fixed by resumming infinite series of diagrams that provide infrared-finite results.

The fields that describe a system of particles at finite temperature in the imaginary time formalism have to satisfy periodic (for bosons) or antiperiodic (for fermions) boundary conditions in the Euclidean time variable. This allows us to expand a generic field $\Phi(x, \tau)$ in terms of its Fourier modes $\phi_n(x)$:

$$\Phi(x, \tau) = T \sum_n \phi_n(x) e^{-i\omega_n \tau}. \quad (1)$$

^aAddress after September 1, 1997: CERN – Theory Division, CH-1211 Geneva 23, Switzerland.

In this expression, ω_n is the so-called *Matsubara frequency* and it is $2n\pi T$ for bosons and $(2n+1)\pi T$ for fermions where n is an integer. We can associate a propagator

$$\frac{1}{\omega_n^2 + p^2} \quad (2)$$

to the n th-mode and ω_n then plays the role of its mass. When $\omega_n = 0$ the Fourier mode is massless and is called static. Otherwise, the Fourier mode has a mass $\omega_n \sim T$ and it is called non-static.

If we decouple the non-static modes what remains is a 3-dimensional effective field theory of the static modes. This procedure is known as *dimensional reduction*^{1,2,3}. The effective theory is therefore described by an effective Lagrangian $\mathcal{L}_{eff}$ which has to be compatible with the symmetries of the original theory. It can be written as $\mathcal{L}_{eff} = \mathcal{L}_R + \delta\mathcal{L}$, where $\mathcal{L}_R$ represents the renormalizable terms of the effective Lagrangian and $\delta\mathcal{L}$ represents an infinite series of non-renormalizable terms. The effective coupling constants are determined by matching physical quantities in the full theory and in the effective theories. It is convenient to introduce an infrared cutoff in both theories before matching. The cutoff is chosen to satisfy $gT \ll \Lambda_{IR} \ll T$ so that the strict perturbation expansion (without resummations) is enough to determine the parameters of the effective theory by matching with the full theory^{4,5}.

2 QCD

In this section we will apply these ideas to compute the free energy of QCD. Our goal is to compute the free energy of QCD up to order g^5 . Therefore, we will systematically ignore higher order contributions.

The thermodynamic properties of QCD matter in thermal equilibrium at temperature T are described by the free energy F_{QCD} which is given by the expression

$$e^{-\beta F_{\text{QCD}}} = Z_{\text{QCD}} = \int \mathcal{D}A_\mu(\mathbf{x}, \tau) \mathcal{D}q \mathcal{D}\bar{q} e^{-\int_0^\beta d\tau \int d^3x \mathcal{L}_{\text{QCD}}}, \quad (3)$$

where Z_{QCD} is the partition function, $\beta = 1/T$, and $\mathcal{L}_{\text{QCD}}$ is the Lagrangian of QCD

$$\mathcal{L}_{\text{QCD}} = \frac{1}{4}(G_{\mu\nu}^a)^2 + \bar{q}\gamma_\mu D_\mu q, \quad (4)$$

where $G_{\mu\nu}^a$ is the strength tensor and D_μ the covariant derivative. The gauge coupling constant is g . Note that the parameters of this theory are T and g and the free energy *density* is given by

$$\mathcal{F} = -\frac{T}{V} \log Z_{\text{QCD}}. \quad (5)$$

By dimensional reduction at high T we obtain the effective theory described by the partition function

$$\mathcal{Z}_{\text{EQCD}} = \int \mathcal{D}A_0(\mathbf{x}) \mathcal{D}A_i(\mathbf{x}) e^{-\int d^3x \mathcal{L}_{\text{EQCD}}} . \quad (6)$$

Here, EQCD stands for *Electrostatic QCD*. The Lagrangian is

$$\mathcal{L}_{\text{EQCD}} = \frac{1}{4}(G_{ij}^a)^2 + \frac{1}{2}(D_i A_0^a)(D_i A_0^a) + \frac{1}{2}m_E^2 A_0^a A_0^a + \frac{1}{8}\lambda_E (A_0^a A_0^a)^2 + \delta\mathcal{L}_{\text{EQCD}} , \quad (7)$$

where, $\delta\mathcal{L}_{\text{EQCD}}$ represents an infinite series of non-renormalizable terms. The effective parameters are the effective gauge coupling constant g_E , the effective mass of the electrostatic field m_E , λ_E , and an infinite number of coupling constants associated with the non-renormalizable terms. In addition, since we intend to compute the free energy, we also have to consider the effective parameter, f_E , associated with the unit operator. All of these effective parameters are functions of g and T . λ_E as well as the parameters of the terms in $\delta\mathcal{L}_{\text{EQCD}}$ can be ignored since they only contribute to the free energy at order larger than g^5 ^{6,4,5}. Note that the quarks have been integrated out because their Matsubara modes have a mass of order T . The free energy of QCD can be written as

$$\mathcal{F} = T \left(f_E - \frac{\log \mathcal{Z}_{\text{EQCD}}}{V} \right) . \quad (8)$$

The parameters for EQCD are determined by matching the strict perturbation expansions for the free energy and the electric screening mass in QCD and EQCD^{4,5}. They can be written as an expansion in powers of g^2 .

$$f_E = T^3 (\text{Series in } g^2) , \quad (9)$$

$$m_E^2 = g^2 T^2 (\text{Series in } g^2) , \quad (10)$$

$$g_E^2 = g^2 T (\text{Series in } g^2) , \quad (11)$$

The mass of the electrostatic field $A_0^a(\mathbf{x})$ is $m_E \sim gT$. Therefore, for processes involving scales much smaller than gT (e.g., of order g^2T), $A_0^a(\mathbf{x})$ decouples. What remains is an effective field theory in 3 dimensions which only depends on the magnetostatic fields $A_i^a(\mathbf{x})$ and it is therefore called *Magnetostatic QCD*. It is described by the partition function

$$\mathcal{Z}_{\text{MQCD}} = \int \mathcal{D}A_i(\mathbf{x}) e^{-\int d^3x \mathcal{L}_{\text{MQCD}}} . \quad (12)$$

where

$$\mathcal{L}_{\text{MQCD}} = \frac{1}{4}(G_{ij}^a)^2 + \delta\mathcal{L}_{\text{MQCD}} . \quad (13)$$

In this expression, the strength tensor now involves the effective coupling constant g_M and $\delta\mathcal{L}_{MQCD}$ contains the infinite series of non-renormalizable terms. In addition, we also have to consider the coefficient f_M of the unit operator. The parameters of MQCD are functions of the parameters of EQCD. The free energy of QCD is now written

$$\mathcal{F} = T \left(f_E + f_M - \frac{\log \mathcal{Z}_{MQCD}}{V} \right). \quad (14)$$

f_M is determined by matching the strict perturbation expansions for the free energy in EQCD and MQCD^{4,5}. It can be written as an expansion in powers of g .

$$f_M = m_E^3 (\text{Series in } g), \quad (15)$$

At lowest order in g , it can also be shown^{4,5} that

$$\frac{\log \mathcal{Z}_{MQCD}}{V} \sim g^6 T^3. \quad (16)$$

Therefore, this term does not contribute to the calculation of the free energy of QCD at order g^5 .

We conclude that the free energy of QCD up to order g^5 is given by

$$\mathcal{F} = T (f_E(g, T) + f_M(m_E, g_E)). \quad (17)$$

Details of the calculations of f_E , m_E , g_E , and f_M and complete results are given in Refs. 4 and 5. The result found in this way for $\mathcal{F}$ is in complete agreement with the one obtained by Kastening and Zhai⁷.

3 Convergence of Perturbation Theory

We have outlined the calculation of the free energy as a perturbation expansion in powers of g up to order g^5 . In this section, we examine the convergence of the series and it is based on Ref. 5. We now ask how small $\alpha_s \equiv g^2/(4\pi)$ must be in order for the perturbation expansion to be well-behaved in the sense that every term of a given order is smaller than all of the terms of lower order. If the series is apparently convergent in this sense, then it can plausibly be used to evaluate the free energy.

We study the expansion the of the free energy of QCD in powers of $\sqrt{\alpha_s}$ at different temperatures $T > T_c \sim 200$ MeV. We choose the renormalization scale to be $\mu = 2\pi T$, which is the mass of the lightest non-static mode. In Table 1 we give the contributions coming from each order in $\sqrt{\alpha_s}$, rescaled so that the leading order term is 1, the next terms are of order α_s , $\alpha_s^{3/2}$, α_s^4 ,

T (GeV)	$\alpha_s(2\pi T)$	expansion for F
0.250	0.321	$1 - 0.282 + 0.583 + 0.276 - 1.094$
0.500	0.239	$1 - 0.210 + 0.374 + 0.010 - 0.524$
1	0.194	$1 - 0.167 + 0.267 + 0.033 - 0.287$
2	0.165	$1 - 0.142 + 0.209 + 0.011 - 0.191$
1000	0.074	$1 - 0.062 + 0.062 + 0.011 - 0.024$

Table 1: Perturbation expansion for the free energy of QCD at different temperatures.

and $\alpha_s^{5/2}$, respectively. We see that the correction of order $\alpha_s^{5/2}$ is the largest unless the temperature T is greater than 2 GeV. Also, the term of order $\alpha_s^{3/2}$ is smaller than the term of order α_s only when the temperature is greater than about 1 TeV.

We can go further in our analysis provided that we have separated the contributions from the scales T and gT . The term f_E gives the contribution to the free energy from the scale T and f_M gives the contribution from the scale gT . The analysis shows that the perturbation series for f_E is well-behaved even at temperatures as low as 250 MeV. On the contrary, the temperature for which the perturbation series of f_M is well-behaved is larger than 2 GeV. This suggests that the slow convergence of the expansion for the free energy of QCD in powers of $\sqrt{\alpha_s}$ may be attributed to the slow convergence of perturbation theory at the scale gT .

Acknowledgments

This work was supported in part by the U.S. Department of Energy, Division of High Energy Physics, under Grant DE-FG02-91-ER40690.

References

1. T. Appelquist and R.D. Pisarski, *Phys. Rev. D* **23**, 2305 (1981).
2. S. Nadkarni, *Phys. Rev. D* **27**, 917 (1983).
3. D.J. Gross, R.D. Pisarski, and L.G. Yaffe, *Rev. Mod. Phys.* **53**, 43 (1981).
4. E. Braaten and A. Nieto, *Phys. Rev. D* **53**, 3437 (1996).
5. A. Nieto, *Int. J. Mod. Phys. A* **12**, 1431 (1997).
6. E. Braaten and A. Nieto, *Phys. Rev. Lett.* **76**, 1417 (1996).
7. B. Kastening and C. Zhai, *Phys. Rev. D* **52**, 7232 (1995).